

CONSENT FOR CERVICAL LAMINECTOMY / LAMINOPLASTY

I have discussed the operative procedure of cervical laminectomy / laminoplasty with the patient. I have explained the operative procedure with its intended benefits of pain relief, halting neurological deterioration, as well as possibly improving function and symptoms related to walking and use of hands.

I have also discussed the risk of potential complications, in particular alluding to:

Failure of symptoms to resolve or continued progression

Spinal cord injury and nerve root injury.

This can result in increased weakness (permanent or temporary) in the arms and legs and/or sensory disturbance with sensory loss or parasthesiae in the limbs and urinary incontinence due to spincteric weakness but the risk of this resulting in a permanent and significant functional neurological disability (paralysis) is very rare, with a risk of less than 1%. This may be due to direct injury or compression from a haematoma.

Dural tear causing CSF leak.

Wound infection

Wrong level surgery : With appropriate intra - operative imaging undertaking the operation at the wrong level should be considered as an extremely rare complication.

Spinal instability / progressive kyphosis

Post - operative urinary retention which may require the insertion of an indwelling urinary catheter for a few days.

Some of these complications may necessitate further surgery.

Signature
Consenting Neurosurgeon

Signature
Consenting Patient

Please print name

Please print name

Date :